

2016 Florida Youth Substance Abuse Survey

Okeechobee Data Tables

Table 1. Major demographic characteristics of surveyed Okeechobee and Florida Statewide, 2016

	Okeechobee		Florida Statewide	
	N	%	N	%
Sex				
Female	450	51.9	31,515	47.9
Male	400	46.1	32,905	50.0
Race/Ethnic group				
African American	68	7.8	14,666	22.3
American Indian	22	2.5	568	0.9
Asian	10	1.2	743	1.1
Hispanic/Latino	266	30.7	13,174	20.0
Native Hawaiian/Pacific Islander	1	0.1	144	0.2
Other/Multiple	143	16.5	7,523	11.4
White, non-Hispanic	342	39.4	28,309	43.0
Age				
10	1	0.1	65	0.1
11	70	8.1	3,856	5.9
12	120	13.8	8,338	12.7
13	133	15.3	9,230	14.0
14	122	14.1	9,454	14.4
15	132	15.2	10,070	15.3
16	143	16.5	9,684	14.7
17	84	9.7	9,348	14.2
18	45	5.2	4,799	7.3
19 or older	10	1.2	618	0.9
Grade				
6th	166	19.1	9,301	14.1
7th	132	15.2	9,215	14.0
8th	115	13.3	9,326	14.2
9th	164	18.9	10,140	15.4
10th	165	19.0	9,834	15.0
11th	58	6.7	9,254	14.1
12th	67	7.7	8,705	13.2
Overall Middle School	413	47.6	27,678	42.1
Overall High School	454	52.4	37,765	57.4
Total	867	100.0	65,776	100.0

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

Table 2. Percentages of Okeechobee and Florida Statewide youth who reported having used various drugs in their lifetimes, 2016

	Okeechobee							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	27.0	46.9	38.8	38.8	28.1	46.2	38.6	21.8	51.4	41.3	37.1	23.6	51.5	39.1
Blacking Out	--	18.7	18.0	18.8	--	17.3	18.7	--	15.9	16.5	15.4	--	16.1	15.9
Cigarettes	13.6	26.1	17.5	23.2	13.5	26.6	20.8	8.0	18.5	14.0	14.1	8.2	18.5	14.1
Vaporizer/E-Cigarette	18.4	35.1	24.7	30.7	20.1	34.1	28.0	14.6	33.9	24.4	27.1	15.7	34.5	25.8
Marijuana or Hashish	12.5	27.4	21.1	20.4	12.8	27.2	21.2	7.0	31.7	21.4	21.3	8.1	31.8	21.3
Synthetic Marijuana	--	5.5	3.0	7.5	--	5.4	5.5	--	4.9	4.8	5.0	--	4.8	4.9
Inhalants	4.8	3.8	5.3	3.5	4.6	4.6	4.2	6.4	4.6	6.1	4.7	6.4	4.7	5.4
Flakka	--	2.3	1.3	3.1	--	2.7	2.3	--	1.0	0.9	1.1	--	1.0	1.0
Club Drugs	0.6	1.8	0.9	1.7	0.6	2.0	1.3	0.7	3.0	1.9	2.2	0.8	2.9	2.1
LSD, PCP or Mushrooms	2.9	4.9	3.4	5.0	3.9	4.3	4.1	1.0	4.8	2.7	3.7	1.1	4.9	3.2
Methamphetamine	0.7	1.2	0.7	1.3	1.0	1.3	1.0	0.5	0.8	0.6	0.8	0.5	0.8	0.7
Cocaine or Crack Cocaine	0.6	3.0	2.6	1.6	1.2	2.8	2.0	0.8	2.5	1.6	1.9	0.8	2.3	1.8
Heroin	0.2	0.9	1.0	0.3	0.5	0.9	0.6	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Depressants	2.1	4.9	4.6	2.9	2.4	5.2	3.7	2.2	6.5	5.3	4.1	2.4	6.6	4.7
Prescription Pain Relievers	4.0	4.4	5.2	3.3	4.2	4.2	4.2	3.3	5.8	5.3	4.2	3.5	5.7	4.8
Prescription Amphetamines	1.5	2.3	1.8	2.3	1.5	2.0	2.0	1.3	4.7	3.3	3.2	1.4	4.6	3.2
Steroids (without a doctor's order)	0.0	0.8	0.1	0.8	0.3	0.8	0.5	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Over-the-Counter Drugs	1.7	3.1	4.1	1.1	1.5	3.3	2.5	3.2	5.3	4.7	4.2	3.4	5.4	4.4
Needle to Inject Illegal Drugs	--	2.5	2.8	1.5	--	2.9	2.5	--	0.8	0.6	0.8	--	0.8	0.8
Any illicit drug	17.0	31.3	26.1	23.7	16.7	32.8	25.3	15.7	36.4	29.0	26.4	16.6	36.6	27.7
Any illicit drug other than marijuana	11.5	15.4	14.9	13.0	12.2	15.2	13.7	12.0	17.7	16.4	14.2	12.3	17.7	15.3
Alcohol only	13.3	19.7	15.3	19.3	14.5	18.9	17.0	12.5	21.0	18.1	17.0	13.4	21.0	17.5
Alcohol or any illicit drug	31.1	51.2	42.1	43.1	31.9	51.9	42.7	27.8	57.2	46.8	43.1	29.6	57.4	44.8
Any illicit drug, but no alcohol	5.0	4.3	3.8	4.6	4.4	5.8	4.6	6.5	6.1	6.0	6.5	6.5	6.1	6.3

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 3. Percentages of Okeechobee and Florida Statewide youth who reported having used various drugs in the past 30 days, 2016

	Okeechobee							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	11.1	24.9	18.8	19.2	12.6	25.7	19.1	8.3	25.5	19.9	17.0	9.1	25.2	18.3
Binge Drinking	6.2	13.7	9.9	10.7	7.1	13.3	10.5	3.2	10.9	7.9	7.7	3.4	10.8	7.7
Cigarettes	1.5	9.9	5.4	6.4	2.2	9.8	6.3	1.4	4.8	3.3	3.5	1.5	4.6	3.4
Vaporizer/E-Cigarette	7.9	16.0	8.9	15.8	10.4	14.3	12.6	5.1	12.8	8.4	10.6	5.5	12.9	9.6
Marijuana or Hashish	5.1	16.0	10.2	13.4	6.5	13.8	11.5	3.2	17.0	10.9	11.5	3.9	16.9	11.2
Synthetic Marijuana	--	2.1	1.2	3.1	--	2.6	2.1	--	1.0	0.9	1.2	--	1.1	1.0
Inhalants	2.2	2.4	3.0	1.8	2.0	3.1	2.3	2.2	1.2	1.9	1.4	2.1	1.2	1.6
Flakka	--	0.9	0.0	1.7	--	1.2	0.9	--	0.5	0.4	0.7	--	0.5	0.5
Club Drugs	0.2	0.4	0.2	0.5	0.2	0.5	0.3	0.3	0.8	0.5	0.6	0.3	0.7	0.6
LSD, PCP or Mushrooms	1.3	1.0	0.7	1.6	1.5	1.0	1.1	0.3	1.4	0.7	1.1	0.3	1.4	0.9
Methamphetamine	0.0	0.7	0.0	0.8	0.3	0.5	0.4	0.3	0.4	0.3	0.4	0.3	0.4	0.4
Cocaine or Crack Cocaine	0.2	1.0	0.2	1.1	0.5	0.9	0.6	0.3	0.8	0.6	0.6	0.3	0.8	0.6
Heroin	0.0	0.5	0.2	0.5	0.3	0.3	0.3	0.1	0.2	0.1	0.2	0.1	0.2	0.2
Depressants	1.1	2.3	1.5	1.8	1.0	3.0	1.8	0.8	2.4	2.0	1.5	0.9	2.5	1.8
Prescription Pain Relievers	3.1	2.4	3.1	2.4	3.2	2.8	2.7	1.6	2.0	2.2	1.5	1.6	2.1	1.8
Prescription Amphetamines	0.8	1.3	0.4	1.8	0.7	1.7	1.1	0.5	1.6	1.1	1.2	0.6	1.7	1.2
Steroids (without a doctor's order)	0.0	0.6	0.2	0.5	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Over-the-Counter Drugs	1.0	0.9	1.8	0.2	0.4	1.5	0.9	1.8	2.1	2.0	1.9	1.8	2.1	2.0
Any illicit drug	8.6	18.5	14.3	14.9	9.2	17.6	14.3	7.5	20.0	15.1	14.2	8.1	20.0	14.7
Any illicit drug other than marijuana	5.8	6.8	7.1	5.8	5.5	8.6	6.4	5.6	7.7	7.3	6.2	5.7	7.7	6.8
Alcohol only	5.0	12.4	9.8	8.2	6.1	12.8	9.3	5.2	13.4	11.1	9.1	5.6	13.3	10.0
Alcohol or any illicit drug	15.1	30.5	24.2	23.6	16.6	30.0	24.0	12.5	32.9	25.8	22.9	13.5	32.9	24.3
Any illicit drug, but no alcohol	4.4	5.8	6.0	4.6	4.3	4.6	5.2	4.5	7.8	6.4	6.3	4.7	8.0	6.4

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Okeechobee, 2006 to 2016

	2006			2008			2010			2012			2014			2016		
	Middle School	High School	Total															
Alcohol	51.5	66.9	60.2	53.8	76.9	66.7	--	--	--	40.9	67.0	55.5	26.2	63.6	46.8	27.0	46.9	38.6
Blacking Out	--	--	--	--	--	--	--	--	--	--	--	--	--	26.8	26.8	--	18.7	18.7
Cigarettes	32.3	45.5	39.8	33.0	44.5	39.4	--	--	--	23.9	41.4	33.7	13.4	36.9	26.4	13.6	26.1	20.8
Vaporizer/E-Cigarette	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	18.4	35.1	28.0
Marijuana or Hashish	13.4	33.0	24.6	14.5	32.0	24.3	--	--	--	12.4	33.1	24.1	9.5	34.9	23.4	12.5	27.4	21.2
Synthetic Marijuana	--	--	--	--	--	--	--	--	--	--	12.3	12.3	--	14.8	14.8	--	5.5	5.5
Inhalants	14.4	7.1	10.3	18.2	12.1	14.8	--	--	--	12.6	6.8	9.4	6.7	6.3	6.5	4.8	3.8	4.2
Flakka	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	2.3	2.3
Club Drugs	--	--	--	3.1	--	3.1	--	--	--	2.7	4.4	3.7	1.9	4.7	3.4	0.6	1.8	1.3
LSD, PCP or Mushrooms	--	--	--	2.7	--	2.7	--	--	--	2.1	4.9	3.7	3.6	6.3	5.1	2.9	4.9	4.1
Methamphetamine	3.6	3.7	3.6	1.0	3.9	2.6	--	--	--	1.8	2.3	2.0	0.5	1.0	0.8	0.7	1.2	1.0
Cocaine or Crack Cocaine	--	--	--	3.5	--	3.5	--	--	--	2.0	4.6	3.5	1.9	6.3	4.3	0.6	3.0	2.0
Heroin	1.0	1.3	1.2	0.9	1.8	1.4	--	--	--	0.6	1.5	1.1	0.0	0.9	0.5	0.2	0.9	0.6
Depressants	3.1	9.6	6.8	5.0	10.3	7.9	--	--	--	3.0	6.4	4.9	2.6	8.7	5.9	2.1	4.9	3.7
Prescription Pain Relievers	3.4	12.2	8.4	6.7	12.3	9.8	--	--	--	4.8	9.5	7.4	3.6	7.9	5.9	4.0	4.4	4.2
Prescription Amphetamines	3.5	9.9	7.1	2.2	6.5	4.6	--	--	--	2.2	5.6	4.1	1.5	2.4	2.0	1.5	2.3	2.0
Steroids (without a doctor's order)	1.8	0.9	1.3	0.5	1.0	0.8	--	--	--	1.8	1.2	1.4	0.2	0.5	0.3	0.0	0.8	0.5
Over-the-Counter Drugs	--	--	--	4.5	--	4.5	--	--	--	5.9	5.6	5.7	5.4	6.1	5.8	1.7	3.1	2.5
Needle to Inject Drugs	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	2.5	2.5
Any illicit drug	26.2	40.2	34.2	29.9	42.3	36.8	--	--	--	25.0	38.9	32.8	16.9	41.4	30.4	17.0	31.3	25.3
Any illicit drug other than marijuana	21.2	22.5	21.9	25.2	29.8	27.8	--	--	--	20.1	22.0	21.2	13.1	23.1	18.6	11.5	15.4	13.7
Alcohol only	28.9	28.8	28.7	27.8	36.0	32.4	--	--	--	22.3	30.5	26.9	13.2	25.5	20.0	13.3	19.7	17.0
Alcohol or any illicit drug	54.7	68.9	62.6	57.8	78.2	69.1	--	--	--	47.1	69.5	59.7	30.1	66.8	50.3	31.1	51.2	42.7
Any illicit drug, but no alcohol	3.6	2.3	2.8	4.1	1.7	2.8	--	--	--	6.3	2.7	4.3	3.9	3.4	3.6	5.0	4.3	4.6

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Okeechobee, 2006 to 2016

	2006			2008			2010			2012			2014			2016		
	Middle School	High School	Total															
Alcohol	25.0	46.3	37.0	27.8	49.3	39.7	--	--	--	22.1	40.3	32.4	13.6	34.8	25.3	11.1	24.9	19.1
Binge Drinking	13.8	29.9	23.0	14.0	30.7	23.3	--	--	--	13.2	23.0	18.7	6.4	19.9	13.9	6.2	13.7	10.5
Cigarettes	8.2	20.5	15.0	7.3	18.4	13.5	--	--	--	4.5	17.1	11.6	2.6	11.2	7.3	1.5	9.9	6.3
Vaporizer/E-Cigarette	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	7.9	16.0	12.6
Marijuana or Hashish	6.5	15.4	11.4	6.6	15.4	11.5	--	--	--	5.1	17.9	12.3	6.7	16.1	11.9	5.1	16.0	11.5
Synthetic Marijuana	--	--	--	--	--	--	--	--	--	--	5.3	5.3	--	1.7	1.7	--	2.1	2.1
Inhalants	4.9	3.4	4.1	6.8	0.3	3.2	--	--	--	4.5	1.8	3.0	3.5	1.3	2.3	2.2	2.4	2.3
Flakka	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	0.9	0.9
Club Drugs	--	--	--	0.7	--	0.7	--	--	--	0.9	1.9	1.5	0.9	0.5	0.7	0.2	0.4	0.3
LSD, PCP or Mushrooms	--	--	--	0.5	--	0.5	--	--	--	0.9	1.5	1.2	1.1	1.0	1.1	1.3	1.0	1.1
Methamphetamine	1.0	1.2	1.1	0.5	0.9	0.7	--	--	--	0.8	0.7	0.7	0.4	0.0	0.2	0.0	0.7	0.4
Cocaine or Crack Cocaine	--	--	--	1.7	--	1.7	--	--	--	0.9	1.8	1.4	0.7	1.7	1.2	0.2	1.0	0.6
Heroin	0.3	0.7	0.5	0.0	0.0	0.0	--	--	--	0.0	0.9	0.5	0.0	0.3	0.2	0.0	0.5	0.3
Depressants	1.0	4.2	2.8	1.8	4.6	3.4	--	--	--	1.5	1.9	1.7	1.1	2.2	1.7	1.1	2.3	1.8
Prescription Pain Relievers	0.8	4.5	2.9	3.3	4.7	4.1	--	--	--	1.4	2.9	2.3	1.9	1.6	1.7	3.1	2.4	2.7
Prescription Amphetamines	1.5	4.1	3.0	0.8	0.4	0.6	--	--	--	1.0	1.4	1.2	0.9	0.4	0.6	0.8	1.3	1.1
Steroids (without a doctor's order)	0.5	0.9	0.7	0.4	0.2	0.3	--	--	--	0.0	0.4	0.2	0.0	0.3	0.1	0.0	0.6	0.3
Over-the-Counter Drugs	--	--	--	2.4	--	2.4	--	--	--	2.3	1.9	2.1	3.0	1.8	2.4	1.0	0.9	0.9
Any illicit drug	11.5	21.4	16.9	15.4	19.1	17.5	--	--	--	11.0	21.3	16.8	9.7	20.5	15.7	8.6	18.5	14.3
Any illicit drug other than marijuana	8.6	11.8	10.4	12.2	10.1	11.0	--	--	--	7.2	8.4	7.9	5.8	7.8	6.9	5.8	6.8	6.4
Alcohol only	17.1	29.3	24.1	17.1	33.2	26.0	--	--	--	14.4	24.1	19.9	7.6	20.9	15.0	5.0	12.4	9.3
Alcohol or any illicit drug	28.5	50.6	41.0	32.1	51.8	43.1	--	--	--	25.9	45.0	36.6	17.2	41.1	30.4	15.1	30.5	24.0
Any illicit drug, but no alcohol	4.0	4.4	4.2	4.7	2.8	3.7	--	--	--	3.7	4.8	4.3	3.9	6.6	5.4	4.4	5.8	5.2

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 6. Percentages of Okeechobee and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, parental disapproval and peer disapproval, 2016

	Okeechobee							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Early ATOD use														
More than a sip of alcohol	--	25.3	24.7	25.6	--	24.8	25.3	--	19.4	18.3	20.2	--	19.3	19.4
Drinking at least once a month	--	5.9	5.6	6.5	--	6.3	5.9	--	3.5	3.7	3.4	--	3.5	3.5
Smoke cigarettes	--	15.9	12.6	19.0	--	17.2	15.9	--	9.3	8.6	9.8	--	9.4	9.3
Smoke marijuana	--	14.0	10.2	17.8	--	13.2	14.0	--	10.6	9.0	12.0	--	10.8	10.6
Great risk of harm														
1 or 2 drinks nearly every day	38.1	36.8	38.1	37.2	38.4	33.7	37.3	45.7	40.7	46.4	39.3	45.9	40.0	42.8
5+ drinks once or twice a week	50.5	44.4	48.6	45.9	50.6	43.0	47.0	58.1	52.3	58.4	51.2	58.4	51.8	54.7
1+ packs of cigarettes per day	60.7	62.4	61.2	61.6	60.9	64.0	61.7	67.9	68.8	68.4	68.5	68.7	68.2	68.4
Marijuana once or twice a week	44.5	30.8	36.6	36.6	44.8	28.8	36.6	51.7	25.2	37.5	35.2	50.1	24.4	36.3
Try marijuana once or twice	32.4	23.1	27.7	26.9	33.1	20.0	27.0	36.1	16.8	24.8	25.0	34.7	16.4	24.9
Use prescription drugs	68.5	66.9	67.9	68.0	69.0	65.9	67.5	69.1	68.2	70.0	67.3	69.4	67.9	68.5
Think it would be wrong to...														
Smoke cigarettes	92.8	82.4	89.6	85.2	91.2	85.1	86.8	95.6	87.7	91.1	91.0	95.3	88.9	91.0
Drink alcohol regularly	83.4	62.6	72.8	70.3	81.9	65.0	71.4	88.8	64.5	74.1	75.2	87.6	64.8	74.7
Smoke marijuana	87.4	65.2	75.6	74.5	85.5	68.2	74.6	89.7	62.3	74.4	73.3	88.1	62.3	73.8
Smoke synthetic marijuana	90.6	89.0	91.0	88.7	89.4	89.9	89.7	93.1	88.6	90.1	90.9	92.7	88.6	90.5
Use other illicit drugs	96.7	92.1	95.6	93.2	95.8	92.4	94.1	97.5	93.6	95.8	94.7	97.3	93.6	95.2
Parents think it would be wrong to...														
1 or 2 drinks nearly every day	96.1	92.1	94.3	93.5	96.7	91.5	93.8	97.3	93.8	95.4	95.1	97.2	94.1	95.2
Smoke cigarettes	98.8	92.7	94.2	96.7	99.1	94.8	95.3	98.6	96.8	97.6	97.5	98.6	97.3	97.6
Smoke marijuana	95.9	86.9	89.9	91.9	95.7	91.0	90.8	97.0	88.6	92.3	92.0	96.6	88.9	92.1
Use prescription drugs	98.0	97.1	97.5	97.4	97.9	98.7	97.5	98.2	97.3	97.4	97.9	98.1	97.4	97.7
Friends think it would be wrong to...														
1 or 2 drinks nearly every day	86.6	71.5	81.4	74.7	86.2	70.4	77.8	91.3	79.0	85.6	82.9	90.7	79.1	84.2
Smoke tobacco	93.4	79.1	87.4	83.6	92.6	79.6	85.2	94.8	87.0	91.1	89.5	94.4	87.8	90.3
Smoke marijuana	81.7	63.1	69.6	72.2	81.3	63.8	71.0	88.6	59.9	72.6	71.3	86.6	59.8	72.0
Use prescription drugs	95.9	91.3	95.4	91.7	95.9	90.1	93.2	95.7	90.6	93.4	92.2	95.5	90.3	92.7

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

Table 7. Trends in early ATOD use and attitudes toward substance use for Okeechobee, 2006 to 2016

	2006			2008			2010			2012			2014			2016		
	Middle School	High School	Total															
Early ATOD use																		
Try alcohol	--	34.6	34.6	--	38.7	38.7	--	--	--	--	31.2	31.2	--	26.8	26.8	--	25.3	25.3
Drink monthly	--	6.2	6.2	--	10.3	10.3	--	--	--	--	10.2	10.2	--	4.9	4.9	--	5.9	5.9
Smoke cigarettes	--	32.4	32.4	--	32.5	32.5	--	--	--	--	19.4	19.4	--	23.0	23.0	--	15.9	15.9
Smoke marijuana	--	17.0	17.0	--	17.6	17.6	--	--	--	--	13.7	13.7	--	15.0	15.0	--	14.0	14.0
Great risk of harm																		
Daily alcohol use	34.6	35.2	34.9	35.9	33.8	34.7	--	--	--	38.7	31.9	34.9	39.3	26.0	31.9	38.1	36.8	37.3
5+ drinks weekly	--	--	--	--	--	--	--	--	--	--	--	--	47.3	41.1	43.8	50.5	44.4	47.0
Daily cigarette use	63.4	53.6	58.0	64.0	65.6	64.9	--	--	--	60.7	59.6	60.0	61.5	59.9	60.6	60.7	62.4	61.7
Marijuana weekly*	--	--	--	--	--	--	--	--	--	--	--	--	47.4	24.4	34.6	44.5	30.8	36.6
Try marijuana	44.0	34.5	38.5	42.4	23.4	31.9	--	--	--	35.5	23.3	28.6	39.1	16.5	26.4	32.4	23.1	27.0
Prescription drugs	--	--	--	--	--	--	--	--	--	--	74.0	74.0	70.6	75.7	73.4	68.5	66.9	67.5
Think it wrong																		
Smoke cigarettes	85.9	65.6	74.5	84.0	60.5	70.8	--	--	--	88.7	70.9	78.6	92.0	79.7	85.1	92.8	82.4	86.8
Drink regularly	72.9	48.7	59.4	70.8	44.6	56.1	--	--	--	75.0	51.7	61.8	85.4	56.9	69.3	83.4	62.6	71.4
Smoke marijuana	89.3	75.0	81.4	87.1	74.2	79.8	--	--	--	88.6	68.5	77.2	87.8	63.9	74.3	87.4	65.2	74.6
Smoke synthetic marijuana	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	90.6	89.0	89.7
Use other illicit drugs	97.2	92.2	94.4	94.4	91.6	92.8	--	--	--	95.6	94.3	94.8	96.1	94.7	95.3	96.7	92.1	94.1
Parents think it wrong																		
Daily alcohol use*	--	--	--	--	--	--	--	--	--	--	--	--	96.2	87.8	91.6	96.1	92.1	93.8
Smoke cigarettes	94.8	87.8	90.6	94.7	86.3	90.1	--	--	--	96.4	91.3	93.5	97.5	94.3	95.8	98.8	92.7	95.3
Smoke marijuana	95.9	90.0	92.4	97.1	93.2	95.0	--	--	--	94.9	92.2	93.4	96.3	87.1	91.3	95.9	86.9	90.8
Prescription drugs	--	--	--	--	--	--	--	--	--	--	--	--	98.4	97.8	98.1	98.0	97.1	97.5
Friends think it wrong																		
Daily alcohol use	--	--	--	--	--	--	--	--	--	--	--	--	88.0	66.5	76.1	86.6	71.5	77.8
Smoke tobacco	--	--	--	--	--	--	--	--	--	--	--	--	92.7	75.9	83.3	93.4	79.1	85.2
Smoke marijuana	--	--	--	--	--	--	--	--	--	--	--	--	88.5	59.9	72.6	81.7	63.1	71.0
Prescription drugs	--	--	--	--	--	--	--	--	--	--	--	--	94.4	92.3	93.2	95.9	91.3	93.2

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

* In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." Also, in 2014, the description of alcohol use in the parent disapproval question was changed from "regularly" to "nearly every day." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among high school drinkers, usual source of alcohol within the past 30 days, Okeechobee and Florida Statewide youth, 2016

	Okeechobee						Florida Statewide							
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Bought in a store	--	5.9	3.3	9.5	--	6.0	5.9	--	8.3	5.0	12.4	--	7.3	8.3
Bought in a restaurant, bar, or club	--	3.8	3.0	0.0	--	5.0	3.8	--	1.6	1.2	2.1	--	1.5	1.6
Bought at a public event	--	2.2	1.9	2.9	--	2.9	2.2	--	0.7	0.7	0.8	--	0.6	0.7
Someone bought it for me	--	16.4	14.3	13.5	--	18.9	16.4	--	14.7	13.3	16.3	--	14.5	14.7
Someone gave it to me	--	45.6	46.8	50.8	--	43.5	45.6	--	44.8	50.6	37.8	--	47.0	44.8
Took it from a store	--	0.6	1.3	0.0	--	0.8	0.6	--	0.3	0.2	0.4	--	0.4	0.3
Took it from a family member	--	5.1	7.0	3.7	--	3.2	5.1	--	11.7	14.2	8.7	--	12.4	11.7
Some other way	--	20.4	22.4	19.6	--	19.7	20.4	--	17.8	14.7	21.5	--	16.5	17.8

Note: The symbol "--" indicates that data are not available.

Table 9. Among high school drinkers, usual drinking location within the past 30 days, Okeechobee and Florida Statewide youth, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
My home	--	36.5	34.2	35.3	--	37.6	36.5	--	37.7	39.2	35.9	--	37.1	37.7
Another person's home	--	30.9	32.2	33.5	--	28.4	30.9	--	40.0	41.0	38.9	--	40.8	40.0
Car or other vehicle	--	0.0	0.0	0.0	--	0.0	0.0	--	1.7	1.6	1.9	--	1.7	1.7
Restaurant, bar, or club	--	1.5	1.9	1.3	--	1.9	1.5	--	2.7	2.9	2.5	--	2.4	2.7
Public place	--	4.1	7.7	1.3	--	2.4	4.1	--	3.9	3.9	3.7	--	4.3	3.9
Public event	--	0.0	0.0	0.0	--	0.0	0.0	--	1.6	1.8	1.5	--	1.7	1.6
School property	--	6.1	4.6	8.2	--	4.9	6.1	--	1.3	0.7	2.0	--	1.4	1.3
Some other place	--	21.0	19.4	20.4	--	24.7	21.0	--	11.1	9.0	13.7	--	10.6	11.1

Note: The symbol "--" indicates that data are not available.

Table 10. Among high school drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Okeechobee and Florida Statewide youth, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
1	--	18.3	26.0	12.3	--	18.1	18.3	--	29.6	31.6	27.4	--	30.5	29.6
2	--	22.4	22.3	25.1	--	23.6	22.4	--	22.5	24.8	19.8	--	22.4	22.5
3	--	16.2	21.2	13.6	--	15.2	16.2	--	17.3	17.9	16.3	--	16.9	17.3
4	--	10.5	16.0	6.6	--	11.0	10.5	--	8.6	9.0	8.3	--	8.5	8.6
5 or more	--	32.6	14.6	42.4	--	32.1	32.6	--	22.0	16.7	28.2	--	21.8	22.0

Note: The symbol "--" indicates that data are not available.

Table 11. Percentages of Okeechobee and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high before or during school in the past 12 months, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Drinking alcohol	4.5	7.9	5.2	7.7	4.6	7.6	6.5	3.3	6.5	5.5	4.7	3.3	6.8	5.1
Smoking marijuana	6.3	10.4	7.3	10.4	6.5	9.2	8.6	3.2	12.2	8.1	8.7	3.7	12.3	8.5
Used another drug to get high	2.1	3.0	1.6	3.7	2.4	3.6	2.6	1.4	3.9	2.9	2.9	1.5	4.1	2.9

Table 12. Percentages of Okeechobee and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Carrying a handgun	10.3	7.8	3.4	13.8	8.7	9.4	8.9	5.0	5.8	2.6	8.3	5.0	5.9	5.5
Selling drugs	4.3	4.8	2.5	6.8	3.8	4.8	4.6	1.6	6.0	3.0	5.4	1.8	6.1	4.2
Attempting to steal a vehicle	1.4	2.3	0.3	3.5	1.7	2.2	1.9	1.0	1.5	0.8	1.7	1.0	1.5	1.3
Being arrested	4.5	3.4	1.4	6.2	4.2	3.5	3.9	1.9	2.8	1.8	3.0	1.8	3.0	2.4
Taking a handgun to school	1.0	1.4	0.4	2.2	1.1	1.2	1.3	0.4	0.7	0.3	0.9	0.4	0.7	0.6
Getting suspended	15.8	13.1	9.4	17.7	15.5	13.4	14.3	10.6	9.1	7.0	12.5	10.1	9.8	9.8
Attacking someone with intent to harm	8.3	5.9	4.7	9.0	6.4	8.4	7.0	6.2	6.1	5.3	6.9	6.0	6.6	6.1

Table 13. Trends in delinquent behaviors for Okeechobee, 2006 to 2016

	2006			2008			2010			2012			2014			2016		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>															
Carrying a handgun	9.3	9.1	9.4	7.4	10.0	8.9	--	--	--	7.4	6.4	6.9	9.8	8.3	9.0	10.3	7.8	8.9
Selling drugs	2.1	6.9	5.0	4.1	10.4	7.6	--	--	--	3.6	6.8	5.4	2.9	6.2	4.7	4.3	4.8	4.6
Attempting to steal a vehicle	2.4	5.8	4.5	4.4	2.4	3.3	--	--	--	1.9	2.6	2.3	2.5	1.3	1.9	1.4	2.3	1.9
Being arrested	6.1	8.3	7.4	5.3	6.0	5.7	--	--	--	3.8	4.7	4.3	3.3	5.0	4.2	4.5	3.4	3.9
Taking a handgun to school	0.6	2.9	2.0	0.8	1.7	1.3	--	--	--	0.3	1.4	0.9	0.2	0.6	0.4	1.0	1.4	1.3
Getting suspended	21.5	23.0	22.3	17.0	24.0	20.8	--	--	--	13.6	16.0	14.9	15.1	10.3	12.5	15.8	13.1	14.3
Attacking someone with intent to harm	13.8	15.1	14.6	15.4	21.2	18.6	--	--	--	11.0	12.8	12.0	7.7	7.6	7.7	8.3	5.9	7.0

Table 14. Percentages of Okeechobee and Florida Statewide youth with family members who have been or are now in jail or friends who have been in trouble because of alcohol or drugs, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Family members who have been or are now in jail or prison	57.6	49.5	59.2	47.4	55.2	51.1	52.9	44.4	44.6	47.8	41.2	44.2	45.4	44.5
Friends who have been or are in trouble because of alcohol or drugs	31.4	44.7	39.6	39.7	29.9	45.8	39.1	23.9	46.5	38.4	35.8	26.0	47.1	37.1

Table 15. Percentages of Okeechobee and Florida Statewide youth who reported involvement in bullying behavior, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Skipped school because of bullying	10.2	9.1	13.9	4.6	10.4	8.1	9.6	7.6	8.2	11.5	4.4	7.6	8.3	7.9
Was kicked or shoved in past 30 days	19.1	7.5	13.4	11.4	18.3	7.7	12.3	19.6	8.1	12.0	13.8	18.8	8.1	13.0
Was taunted or teased in past 30 days	35.9	16.0	31.2	17.7	34.6	17.1	24.4	35.3	22.5	30.1	25.6	34.4	22.8	27.9
Was victim of cyber bullying in past 30 days	12.4	7.3	13.4	6.1	12.6	6.6	9.4	7.8	7.6	10.2	5.2	7.8	7.9	7.7
Physically bullied others in past 30 days	5.9	5.1	4.6	6.4	6.0	5.3	5.4	7.6	4.9	5.0	7.0	7.2	5.2	6.1
Verbally bullied others in past 30 days	15.7	8.0	10.7	12.1	15.2	7.7	11.3	14.3	10.4	11.0	13.0	14.0	10.7	12.0
Cyber bullied others in past 30 days	5.1	3.1	4.6	3.4	5.1	2.5	3.9	3.6	4.1	4.0	3.7	3.7	4.1	3.9

Table 16. Percentages of Okeechobee and Florida Statewide youth who reported gang membership, 2016

	Okeechobee							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Have you ever belonged to a gang?	4.4	5.5	2.7	7.6	3.7	7.3	5.0	3.3	3.5	2.3	4.5	3.1	3.6	3.4
Did that gang have a name?	28.3	11.2	9.2	19.6	20.2	15.2	14.2	23.4	14.1	13.1	19.7	21.9	14.4	16.9
Are you a gang member now?	--	3.4	2.1	4.8	--	4.2	3.4	--	2.0	1.1	2.8	--	2.0	2.0

Note: The prevalence rates for “Did that gang have a name?” exclude students who reported that they have never belonged to a gang. The symbol “--” indicates that data are not available.

Table 17. Reasons for joining a gang, reported by high school students who have belonged to a gang, among Okeechobee and Florida Statewide youth, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Fun and excitement	--	4.6	3.7	5.3	--	6.0	4.6	--	5.0	4.3	5.5	--	5.0	5.0
Protection	--	4.7	2.5	6.4	--	5.2	4.7	--	4.2	2.6	5.4	--	4.5	4.2
Friend or relative in a gang	--	8.2	1.5	13.2	--	9.3	8.2	--	4.1	2.9	4.9	--	4.1	4.1
Forced to join	--	0.0	0.0	0.0	--	0.0	0.0	--	1.1	1.1	1.2	--	1.2	1.1
To get respect	--	5.5	5.1	5.8	--	7.1	5.5	--	3.5	2.0	4.7	--	3.5	3.5
Money	--	3.3	1.4	4.7	--	4.2	3.3	--	3.8	2.5	4.8	--	4.1	3.8
To fit in better	--	1.6	2.4	1.1	--	2.1	1.6	--	1.2	1.1	1.4	--	1.2	1.2
Other reasons	--	8.1	7.5	8.7	--	8.6	8.1	--	5.9	4.5	7.0	--	5.6	5.9

Note: The symbol "--" indicates that data are not available.

Table 18. Percentages of Okeechobee and Florida Statewide youth who reported participation in extracurricular activities, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
School Sports	30.9	39.4	29.6	42.0	32.1	39.3	35.8	36.0	40.4	34.5	42.5	36.4	41.4	38.5
Organized Sports Outside of School	37.3	16.3	27.6	23.4	34.5	19.2	25.2	41.6	22.4	28.2	32.9	40.1	22.8	30.5
School Band	22.7	10.8	19.5	12.0	22.0	10.2	15.8	16.7	8.1	12.3	11.1	16.2	8.1	11.7
School Club(s)	20.7	18.4	24.5	15.4	20.1	19.2	19.4	22.1	31.8	35.4	20.4	22.8	31.6	27.6
Community Club(s)	11.5	15.5	17.2	10.7	11.5	13.4	13.8	9.3	12.8	14.6	8.2	9.4	12.6	11.3

Table 19. Percentages of high school students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Okeechobee and Florida Statewide youth, 2016

	Okeechobee							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
<i>Riding in a vehicle</i>														
Alcohol	--	20.8	19.7	22.0	--	19.4	20.8	--	16.4	17.5	15.3	--	16.3	16.4
Marijuana	--	24.7	27.4	22.0	--	24.2	24.7	--	22.7	23.7	21.7	--	22.9	22.7
<i>Driving a vehicle</i>														
Alcohol	--	9.8	7.0	11.9	--	8.3	9.8	--	5.4	5.0	5.8	--	5.2	5.4
Marijuana	--	12.9	11.4	14.3	--	11.7	12.9	--	10.3	9.1	11.4	--	9.9	10.3

Note: The symbol "--" indicates that data are not available.

Table 20. Trends for high school students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for Okeechobee, 2012 and 2016

	2006			2008			2010			2012			2014			2016		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>															
<i>Riding in a vehicle</i>																		
Alcohol	--	--	--	--	--	--	--	--	--	--	30.6	30.6	--	27.8	27.8	--	20.8	20.8
Marijuana	--	--	--	--	--	--	--	--	--	--	24.6	24.6	--	26.3	26.3	--	24.7	24.7
<i>Driving a vehicle</i>																		
Alcohol	--	--	--	--	--	--	--	--	--	--	11.5	11.5	--	11.4	11.4	--	9.8	9.8
Marijuana	--	--	--	--	--	--	--	--	--	--	10.5	10.5	--	12.7	12.7	--	12.9	12.9

Note: The symbol "--" indicates that data are not available.

Table 21. Protective factor prevalence rates for Okeechobee, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Okeechobee		Florida Statewide		National Norms	
		<i>Middle School</i>	<i>High School</i>	<i>Middle School</i>	<i>High School</i>	<i>Middle School</i>	<i>High School</i>
Community	Community Rewards for Prosocial Involvement	45	68	46	69	56	63
Family	Family Opportunities for Prosocial Involvement	55	59	60	59	59	54
	Family Rewards for Prosocial Involvement	52	56	56	56	54	55
School	School Opportunities for Prosocial Involvement	45	54	53	63	57	60
	School Rewards for Prosocial Involvement	53	55	49	59	53	58
Peer and Individual	Religiosity	48	58	49	57	56	62
Average Prevalence Rate		50	58	52	61	56	59

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 22. Risk factor prevalence rates for Okeechobee, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Okeechobee		Florida Statewide		National Norms	
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Disorganization	54	48	42	44	47	47
	Transitions and Mobility	55	57	59	61	47	46
	Laws and Norms Favorable to Drug Use	45	34	37	31	42	42
	Perceived Availability of Drugs	38	27	37	27	45	45
	Perceived Availability of Handguns	28	41	24	36	25	42
Family	Poor Family Management	49	44	40	38	44	45
	Family Conflict	38	30	38	33	42	37
School	Poor Academic Performance	57	51	42	44	45	48
	Lack of Commitment to School	61	63	53	54	47	46
Peer and Individual	Favorable Attitudes toward Antisocial Behavior	45	36	39	35	40	46
	Favorable Attitudes toward ATOD Use	39	37	32	36	39	45
	Early Initiation of Drug Use	28	29	23	22	41	46
Average Prevalence Rate		45	41	39	38	42	45

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 23. Protective factor prevalence rate trends among middle school students for Okeechobee, 2006 to 2016

Domain	Scale	Okeechobee					
		2006	2008	2010	2012	2014	2016
Community	Community Rewards for Prosocial Involvement	48	54	--	54	48	45
Family	Family Opportunities for Prosocial Involvement	58	52	--	55	52	55
	Family Rewards for Prosocial Involvement	50	44	--	48	41	52
School	School Opportunities for Prosocial Involvement	44	44	--	48	50	45
	School Rewards for Prosocial Involvement	40	41	--	52	44	53
Peer and Individual	Religiosity	51	52	--	54	52	48
Average Prevalence Rate		49	48	--	52	48	50

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 24. Risk factor prevalence rate trends among middle school students for Okeechobee, 2006 to 2016

Domain	Scale	Okeechobee					
		2006	2008	2010	2012	2014	2016
Community	Community Disorganization	53	60	--	58	56	54
	Transitions and Mobility	58	59	--	58	52	55
	Laws and Norms Favorable to Drug Use	54	57	--	45	44	45
	Perceived Availability of Drugs	50	63	--	44	40	38
	Perceived Availability of Handguns	40	46	--	37	34	28
Family	Poor Family Management	53	54	--	49	47	49
	Family Conflict	46	51	--	42	38	38
School	Poor Academic Performance	53	51	--	48	49	57
	Lack of Commitment to School	58	58	--	53	55	61
Peer and Individual	Favorable Attitudes toward Antisocial Behavior	59	55	--	49	36	45
	Favorable Attitudes toward ATOD Use	55	53	--	48	32	39
	Early Initiation of Drug Use	55	52	--	43	25	28
Average Prevalence Rate		54	52	--	49	42	45

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 25. Protective factor prevalence rate trends among high school students for Okeechobee, 2006 to 2016

Domain	Scale	Okeechobee					
		2006	2008	2010	2012	2014	2016
Community	Community Rewards for Prosocial Involvement	58	55	--	59	60	68
Family	Family Opportunities for Prosocial Involvement	46	50	--	54	58	59
	Family Rewards for Prosocial Involvement	47	49	--	53	52	56
School	School Opportunities for Prosocial Involvement	52	60	--	52	56	54
	School Rewards for Prosocial Involvement	48	57	--	51	50	55
Peer and Individual	Religiosity	60	61	--	62	60	58
Average Prevalence Rate		52	55	--	55	56	58

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 26. Risk factor prevalence rate trends among high school students for Okeechobee, 2006 to 2016

Domain	Scale	Okeechobee					
		2006	2008	2010	2012	2014	2016
Community	Community Disorganization	50	56	--	56	54	48
	Transitions and Mobility	64	72	--	59	63	57
	Laws and Norms Favorable to Drug Use	45	50	--	47	41	34
	Perceived Availability of Drugs	45	54	--	43	36	27
	Perceived Availability of Handguns	52	58	--	51	56	41
Family	Poor Family Management	57	53	--	44	41	44
	Family Conflict	35	41	--	35	35	30
School	Poor Academic Performance	50	52	--	50	49	51
	Lack of Commitment to School	51	50	--	48	59	63
Peer and Individual	Favorable Attitudes toward Antisocial Behavior	53	55	--	42	42	36
	Favorable Attitudes toward ATOD Use	43	50	--	43	43	37
	Early Initiation of Drug Use	41	46	--	35	35	29
Average Prevalence Rate		45	49	--	46	46	41

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.